

ENDODONTICS



Colleagues for Excellence

PUBLISHED FOR THE DENTAL PROFESSIONAL COMMUNITY BY THE AMERICAN ASSOCIATION OF ENDODONTISTS

Spring/Summer 2004

Restoration of Endodontically Treated Teeth: *The Endodontist's Perspective, Part 1*

Welcome to *ENDODONTICS: Colleagues for Excellence*...the newsletter covering the latest in endodontic treatment, research and technology. We hope you enjoy our coverage on the full scope of options available for patients through endodontic treatment and that you find this information valuable in your practice. Future issues of *ENDODONTICS* will keep you up to date on the state of the art and science in endodontic treatment.

How best to restore teeth after root canal treatment has long been a subject of debate and remains controversial to this day. Volumes of literature exist on the topic, much of it contradictory. Due to our limited space in this publication, we cannot complete an exhaustive review of the issues but will address those of primary importance from the endodontist's perspective. This issue is the first of a two part series, with Part II focusing on disassembly of a restored endodontically treated tooth.

Before Root Canal Treatment

The best plan for success is to begin with the end in mind. Before initiating treatment, the practitioner should carefully examine the tooth for caries and fracture. The tooth should be assessed for restorability, occlusal function and periodontal health, and issues such as biological width and crown-to-root ratio should be evaluated. If these factors are deemed satisfactory, the tooth can be included in the comprehensive treatment plan.

Whenever possible, practitioners should remove all existing restorations and caries before initiating root canal treatment. This allows more accurate assessment of restorability and evaluation for fractures. Teeth with extensive destruction of tooth structure may need crown lengthening or orthodontic eruption prior to endodontic treatment.

The Root Canal Treatment Is Complete — Now What?

Preventing contamination of the root canal system between completion of endodontic treatment and restoration of the tooth

should be a primary concern. Contamination is thought to be an important cause of future problems in endodontically treated teeth. Therefore, an immediate restoration should be placed, whenever possible. Delaying restorative treatment to assess the success of endodontic treatment is generally not in the best interest of the patient. Temporary restorations do not effectively prevent contamination for extended time periods.

When immediate restoration is not possible, orifice barriers should be placed to help protect the root canal system from saliva contamination. Bonded materials such as composite resin or glass ionomer cements are excellent choices. A traditional temporary material such as IRM or Cavit can be used in the access opening. However, the practitioner should be aware that the temporary restoration does not protect the tooth against fracture.

Sometimes endodontic treatment is performed through an existing crown. If the crown appears clinically acceptable, the access opening should be examined for dental caries. Dental caries detector substances can assist in the evaluation, as can magnification. Absence of caries should be assured before the access opening is restored. If caries is present, the first choice is to remove the crown and remove the caries. If this is not practical, remove most of the caries and temporize the tooth as described above. The restorative dentist should then remove the crown and the remaining caries as soon as possible to minimize the chances of contamination of the root canal system.

Temporization of a post space presents a particular problem. Two recent studies showed that a temporary post and crown

provided no more protection from contamination than controls without temporary restorations. If it is not possible to perform a post and core buildup immediately and a temporary post and crown are necessary for esthetic reasons, place a barrier over the gutta-percha at the base of the post space. The canal space should be maintained in a bacteria free environment. Saliva, which contains bacteria, should not come in contact with the canal. Use of saliva as a lubricant in post fabrication is not advised, and a rubber dam should be used during the restorative phase.

Basic Principles in the Restoration of Endodontically Treated Teeth

The preponderance of literature supports the following guiding principles:

- *Posterior teeth with root canal treatment should receive cuspal coverage restorations. Bonded restorations, once thought to obviate the need for cuspal coverage, provide only short-term strengthening of the teeth, according to recent studies.*
- *Anterior teeth with minimal loss of tooth structure can be restored conservatively with bonded restorations.*
- *Preservation of coronal and radicular tooth structure is desirable.*
- *The purpose of a post is to retain core buildup.*
- *A ferrule is highly desirable when a post is used. An adequate ferrule is considered a minimum of 2 mm of vertical height and 1 mm of dentin thickness.*

Posts

The purpose of a post is to retain a core that is needed because of extensive loss of coronal tooth structure. Practitioners should avoid using posts when other anatomic features are available to retain the core. Molars may not require posts because a core can usually be retained by the pulp chamber and canals.

When a post is necessary, it should be placed in a distal canal in mandibular molars and the palatal canal in maxillary molars, because the other canals tend to be thinner and more curved. Multiple posts are seldom indicated. Anterior teeth with extensive loss of coronal tooth structure usually need a post because the pulp chamber and single canal are generally not adequate to retain a core. In addition, anterior teeth are subject to lateral forces during function, whereas posterior teeth are subject primarily to vertical forces. Premolars require clinical judgment because of their transitional morphology. The remaining dentinal structure will dictate whether a post is indicated to compensate for extensive loss of coronal tooth structure.

The use of a post carries with it a certain attendant risk of root fracture, particularly if sound dentin is removed during preparation. Preparation of the post space can also lead to perforation apically or a lateral strip perforation in the fluted portion of the mid-root. To avoid these problems, the practitioner who performed the endodontic treatment should prepare the post space. Several studies have shown that immediate creation of the post space may result in a better apical seal, although several studies also reported no difference.

Some of the areas of general agreement regarding post design include:

- When a post is needed, remove little if any additional dentin beyond what is needed to perform the root canal treatment.
- Retain a minimum of 4 mm of gutta-percha apically.
- Use a post designed to incorporate mechanical features that resist rotational forces
- Since forces concentrate at the crest of bone during function, place the post to extend apical to the crest of the bone. One “rule of thumb” is that the post should extend “into bone” at least as far as it protrudes “out of bone.”

Should endodontic retreatment be necessary, “retrievability” is an important property of a post and should be considered in post selection.

Post Materials

Until recent years, nearly all posts were made of metal. Now they are also available in ceramic, composite and fiber-reinforced materials. Every post material has some advantages that can be cited to justify its use. However, every material also has disadvantages.

Cast metal posts

For many years, custom cast metal posts were considered the standard. Although they are still used in some circles, they require more chair time than other posts and involve laboratory procedures and fees. In comparison studies, they generally do not perform as well as other types of posts. For these reasons, they are no longer widely used. Another disadvantage of cast metal posts is that they require temporization. The added procedures, time and need for temporization increase the possibility of contamination of the root canal system. On the positive side, there are several long-term clinical studies that report high success rates with cast gold posts. From the endodontist's perspective, they are easily removed for retreatment.

Prefabricated metal posts

These posts have been widely used for the past 20 years. They can be placed easily and quickly, and a core can be added and prepared at the same appointment. They are available in various metal alloys, some of which are quite strong and allow for placement of a relatively thin post. They can usually be removed if retreatment of the root canal is necessary.

Most of the metal alloys contained in prefabricated metal posts are considered acceptable, with the exception of titanium alloys. Titanium alloys are generally weak and therefore not suitable for thin posts. In addition, post removal devices are often ineffective because of the softness of the metal. Titanium alloys have the same radiodensity as gutta-percha and are sometimes hard to detect radiographically.

Prefabricated posts are available in active or passive forms. Passive posts are recommended in most cases, but there are a few indications for active posts, primarily in short teeth where retention is minimal. Because active posts have greater potential to cause root fractures and are more difficult to remove, passive posts are therefore preferred for most clinical situations.

Ceramic, glass and zirconium posts

These materials have gained popularity because they are tooth-colored and avoid esthetic problems in the anterior teeth. The only way to remove nonmetallic posts is to grind them out with a bur, a tedious and dangerous procedure. Attachment of the core to the post can also be a problem with the zirconium posts. These posts should be avoided because, if retreatment is needed, it may not be possible, leaving surgery as the next option.

Fiber posts

Most fiber posts contain either carbon fiber or quartz fiber. They have a modulus of elasticity similar to dentin, which allows them to flex with the root when under stress. This is believed to distribute the stresses more evenly throughout the tooth than metal posts, making the root less susceptible to fracture. Some studies have shown that fiber posts strengthen the root when used with a resin luting cement, and several short-term clinical studies have reported high success rates.

The primary concern about fiber posts is whether they allow movement of the core during function or parafunction. If a post has the same modulus of elasticity as the root, but is much thinner in diameter, it will flex more under a load. This may cause leakage under the crown and buildup. Studies are currently underway to address this question and may lead to some reengineering of the current fiber posts. Any initial strengthening of the root by fiber posts is probably lost with time and function.

Carbon fiber posts are generally removed fairly easily by boring through the middle of the post with an ultrasonic or rotary instrument. The alignment of the fibers assists in keeping the instrument moving in the right direction and thus helps prevent perforations. Quartz fiber posts are newer to the market, and there is some disagreement on their ease of removal.

Several short term clinical studies report high success rates with fiber posts. As recalls continue we will be able to further assess their clinical efficacy.

Luting Materials

It is generally accepted that bonded posts are more retentive than cemented posts and provide a better seal. Most clinicians report that a bonded metal post is more difficult to remove than a cemented post. Surprisingly, this is not clearly shown in the literature. A bonded metal post is a double-edged sword. It is possible to get a better seal and perhaps better retention with a bonded post, but retrieval is more difficult. Fiber posts are designed to be used with resin cements, and their ease of removal is not affected by the cement. Generally speaking, if a post has adequate length and resistance form, satisfactory retention and seal can probably be attained with any of the luting agents and post systems.

Conclusion

Practitioners performing endodontic treatment should follow these principles when planning and performing the restoration of endodontically treated teeth:

- Preserve coronal and radicular dentin.
- Avoid contamination of the root canal system.
- Restore the tooth immediately after root canal treatment, if possible.
- Use posts only when necessary to retain a core buildup.
- Restore teeth in a way that allows for future retreatment of the root canal system.
- In most cases, the particular post system used is not as important as following the principles of adequate length, adequate resistance form, adequate strength to allow preservation of dentin, and an adequate ferrule. If these principles are followed, most post systems will perform well.

Everything we do as dentists is temporary with the exception of extractions. We perform procedures with the idea that they will be durable and long lasting, but none of them are “permanent.” Our treatment planning processes should reflect this reality.

Because many important and integrated concepts have been addressed in this issue of ENDODONTICS: Colleagues for Excellence, the AAE encourages readers to review the enclosed reading list to obtain further information.

The AAE Public and Professional Affairs Committee and the Board of Directors developed this issue with special thanks to the author, Dr. Richard S. Schwartz, and prosthodontist Dr. Richard Jordan for his valuable insights.

The information in this newsletter is designed to aid dentists. Practitioners must use their best professional judgment, taking into account the needs of each individual patient when making diagnoses/treatment plans. The AAE neither expressly nor implicitly warrants any positive results, nor expressly nor implicitly warrants against any negative results, associated with the application of this information. If you would like more information, call your endodontic colleague or contact the AAE.

Did you enjoy this issue of *ENDODONTICS*? Did the information have a positive impact on your practice? Are there topics you would like *ENDODONTICS* to cover in the future? We want to hear from you! Send your comments and questions to the American Association of Endodontists at the address below.



ENDODONTICS: *Colleagues for Excellence*

American Association of Endodontists
211 E. Chicago Ave., Suite 1100
Chicago, IL 60611-2691
www.aae.org

Reading List

ENDODONTICS: COLLEAGUES FOR EXCELLENCE, SPRING/SUMMER 2004



Restoration of Endodontically Treated Teeth: The Endodontist's Perspective, Part 1

Prognosis After Restoration of Endodontically Treated Teeth

Sorensen JA, Martinoff JT. Intracoronar reinforcement and coronal coverage: a study of endodontically treated teeth. *J Prosthet Dent* 51:780–784, 1984.

Fredriksson M, Astback J, Pamenius M, et al. A retrospective study of 236 patients with teeth restored by carbon fiber–reinforced epoxy resin posts. *J Prosthet Dent* 80:151–157, 1998.

Ferrari M, Vichi A, Mannocci F, et al. Retrospective study of the clinical performance of fiber posts. *Am J Dent* 13:9B–13B, 2000.

Ferrari M, Vichi A, Garcia-Godoy F, et al. Clinical evaluation of fiber-reinforced epoxy resin posts and cast post and cores. *Am J Dent* 13:15B–18B, 2000.

Aquilino SA, Caplan DJ. Relationship between crown placement and the survival of endodontically treated teeth. *J Prosthet Dent. Mar*; 87(3):256–63, 2002.

Mannocci F, Bertelli E, Sherriff M, Watson TF, Ford TR. Three-year clinical comparison of survival of endodontically treated teeth restored with either full cast coverage or with direct composite restoration. *J Prosthet Dent. Sep*; 88(3):297–301, 2002.

Cheung GS, Chan TK. Long-term survival of primary root canal treatment carried out in a dental teaching hospital. *Int Endod J*; 36:117–28, 2003.

Malferrari S, Monaco C, Scotti R. Clinical evaluation of teeth restored with quartz fiber–reinforced epoxy resin posts. *Int J Prosthodont. Jan–Feb*; 16(1):39–44, 2003.

Iqbal MK, Johansson AA, Akeel RF, Bergenholtz A, Omar R. A retrospective analysis of factors associated with the periapical status of restored, endodontically treated teeth. *Int J Prosthodont. Jan–Feb*; 16(1):31–8, 2003

Indications for a Post

Robbins JW. Guidelines for the restoration of endodontically treated teeth. *J Am Dent Assoc* 120:558–566, 1990.

Kane J, Burgess JO. Modification of the resistance form of amalgam coronal-radicular restorations. *J Prosthet Dent* 65:470–474, 1991.

Baratieri LN, Calderia de Androda MA, Arcar GM, et al. Influence of post placement in the fracture resistance of endodontically treated incisors with direct composite. *J Prosthet Dent* 84 (2):180–184, 2000.

Resistance and Retention Form for Posts

Standlee JP, Caputo AA, Hanson EC. Retention of endodontic dowels: effects of cement, dowel length, diameter, and design. *J Prosthet Dent* 39:401–405, 1978.

Johnson JK, Sakamura JS. Dowel form and tensile force. *J Prosthet Dent* 40:645–649, 1978.

Cooney JP, Caputo AA, Trabert KC. Retention and stress distribution of tapered-end endodontic posts. *J Prosthet Dent* 55:540–546, 1986.

Lambjerg-Hansen H, Asmussen E. Mechanical properties of endodontic posts. *J Oral Rehab* 24:882–87, 1997.

Gallo JR 3rd, Miller T, Xu X, Burgess JO. *In vitro* evaluation of the retention of composite fiber and stainless steel posts. *J Prosthodont. Mar*; 11(1):25–9, 2002.

Al-Wahadni A, Gutteridge DL. An *in vitro* investigation into the effects of retained coronal dentine on the strength of a tooth restored with a cemented post and partial core restoration. *Int Endod J. Nov*; 35(11):913–8, 2002.

Gallo JR 3rd, Miller T, Xu X, Burgess JO. *In vitro* evaluation of the retention of composite fiber and stainless steel posts. *J Prosthodont. Mar*; 11(1):25–9, 2002.

The Ferrule Effect

Sorensen JA, Engelman MJ. Ferrule design and fracture resistance of endodontically treated teeth. *J Prosthet Dent. May*; 63(5):529–36, 1990.

Libman W, Nicholls J. Load fatigue of teeth restored with cast posts and cores and complete crowns. *Int J Prosthet* 1:155–161, 1995.

Isador F, Brondum K, Ravnholt G. The influence of post length and crown ferrule length on the resistance to cyclic loading of bovine teeth with prefabricated titanium posts. *Int J Pros* 12:78–82, 1999.

al-Hazaimah N, Gutteridge DL. An *in vitro* study into the effect of the ferrule preparation on the fracture resistance of crowned teeth incorporating prefabricated post and composite core restorations. *Int Endod J. Jan*; 34(1):40–6, 2001.

Mezzomo E, Massa F, Libera SD. Fracture resistance of teeth restored with two different post-and-core designs cemented with two different cements: an *in vitro* study. Part I. *Quintessence Int. Apr*; 34(4):301–6, 2003.

Zhi-Yue L, Yu-Xing Z. Effects of post-core design and ferrule on fracture resistance of endodontically treated maxillary central incisors. *J Prosthet Dent. Apr*; 89(4):368–73, 2003.

Luting Cements

Duncan JP, Pameijer CH. Retention of parallel-sided titanium posts cemented with six luting agents; an *in vitro* study. *J Prosthet Dent* 80:423–428, 1998.

Ferrari M, Vichi A, Grandini S, Goracci C. Efficacy of a self-curing adhesive-resin cement system on luting glass-fiber posts into root canals: an SEM investigation. *Int J Prosthodont. 14(6)*:543–9, 2001.

Nissan J, Dmitry Y, Assif D. The use of reinforced composite resin cement as compensation for reduced post length. *J Prosthet Dent. Sep*; 86(3):304–8, 2001.

Passive vs. Active Metal Posts

Burns DA, Krause WR, Douglas HB, Burns DR. Stress distribution surrounding endodontic posts. J Prosthet Dent. Oct; 64 (4):412-8, 1990

Felton DA, Webb EL, Kanoy BE, Dugoni J. Threaded endodontic dowels: effects of post design on incidence of root fracture. J Prosthet Dent 65:179-187, 1991.

Standlee JP, Caputo AA. The retentive and stress distributing properties of split threaded endodontic dowels. J Prosthet Dent 68:436-42, 1992.

Burgess JO, Summitt JB, Robbins JW. The resistance to tensile, compression, and torsional forces provided by four post systems. J Prosthet Dent 68:899-903, 1992.

Thorsteinsson TS, Yaman P, Craig RG. Stress analyses of four prefabricated posts. J Prosthet Dent 67:30-33, 1992.

Metal vs. Non-metal Posts

Purton DG, Love RM. Rigidity and retention of carbon fibre versus stainless steel canal posts. Int Endod J 29:262-265, 1996.

Martinez-Insua A, Da Silva L, Rilo B, et al. Comparison of the fracture resistance of pulpless teeth restored with a cast post and core or carbon-fiber post with a composite core. J Prosthet Dent 80:5527-532, 1998.

Mannocci F, Ferrari M, Watson TF. Intermittent loading of teeth restored using quartz fiber, carbon-quartz fiber, and zirconium dioxide ceramic root canal posts. J Adhes Dent. Summer; 1(2):153-8, 1999.

Ottl P, Hahn L, Lauer HC, Fay M. Fracture characteristics of carbon fibre, ceramic and non-palladium endodontic post systems at monotonously increasing loads. J Oral Rehabil. Feb; 29(2):175-83, 2002.

Akkayan B, Gulmez T. Resistance to fracture of endodontically treated teeth restored with different post systems. J Prosthet Dent. Apr; 87(4):431-7, 2002.

Reid LC, Kazemi RB, Meiers JC. Effect of fatigue testing on core integrity and post microleakage of teeth restored with different post systems. J Endod. Feb; 29(2):125-31, 2003.

Coronal Leakage Associated With Post Systems

Bachicha WS, DiFiore PM, Miller DA, Lautenschlager EP, Pashley DH. Microleakage of endodontically treated teeth restored with posts. J Endod. Nov; 24(11):703-8, 1998.

Mannocci F, Ferrari M, Watson TF. Microleakage of endodontically treated teeth restored with fiber posts and composite cores after cyclic loading: a confocal microscopic study. J Prosthet Dent 85:284-91, 2001.

Howdle MD, Fox K, Youngson CC. An *in vitro* study of coronal microleakage around bonded amalgam coronal-radicular cores in endodontically treated molar teeth. Quintessence Int. Jan; 33(1):22-9, 2002.

Core Buildups and Materials

Kovarik RE, Breeding LC, Caughman WF. Fatigue life of three core materials under simulated chewing conditions. J Prosthet Dent 68:584-590, 1992.

Chang W, Millstein P. Effect of design of prefabricated post heads on core materials. J Prosthet Dent 69:475-82, 1993.

Gateau P, Sabek M, Dailey B. Fatigue testing & microscopic evaluation of post & core restorations under artificial crowns. J Prosthet Dent 82:341-347, 1999.

Gateau P, Sabek M, Dailey B. *In vitro* fatigue resistance of glass ionomer cements used in post-and-core applications. J Prosthet Dent. Aug; 86(2):149-55, 2001.

Cohen BI, Pagnillo MK, Newmann I, et al. Retention of a core material supported by three post head designs. J Prosthet Dent 83(6): 624, 2000.

Pilo R, Cardash HS, Levin E, Assif D. Effect of core stiffness on the *in vitro* fracture of crowned, endodontically treated teeth. J Prosthet Dent. Sep; 88(3):302-6, 2002.

Mollersten L, Lockowandt P, Linden LA. A comparison of strengths of five core and post-and-core systems. Quintessence Int. Feb; 33(2):140-9, 2002.

Post Preparations

Kwan EH, Harrington GW. The effect of immediate post preparation of apical seal. J Endod 7:3325-329, 1981.

Dickey DJ, Harris GZ, Lemon RR, Leubke RG. Effect of post space preparation on apical seal using solvent techniques and Peeso reamers. J Endod 8:351-354, 1982.

Fan B, Wu MK, Wesselink PR. Coronal leakage along apical root fillings after immediate and delayed post spaces preparation. Endo Dent Trauma 15:124-7, 1999.

Raiden G, Costa L, Koss S, et al. Residual Thickness of Root in First Maxillary Premolars with Post Space Preparation. J of Endod 25(7):502-505, 1999.

Bonding to Radicular Dentin

Ferrari M, Mannocci F, Vichi A, Cagidiaco MC, Mjor IA. Bonding to root canal: structural characteristics of the substrate. Am J Dent. Oct; 13(5):255-60, 2000.

Mjor IA, Smith MR, Ferrari M, Mannocci F. The structure of dentine in the apical region of human teeth. Int Endod J. Jul; 34(5):346-53, 2001.

Vichi A, Grandini S, Ferrari M. Comparison between two clinical procedures for bonding fiber posts into a root canal: a microscopic investigation. J Endod. May; 28(5):355-60, 2002.

Vichi A, Grandini S, Davidson CL, Ferrari M. An SEM evaluation of several adhesive systems used for bonding fiber posts under clinical conditions. Dent Mater. Nov; 18(7):495-502, 2002.

Kijsamanmith K, Timpawat S, Harnirattisai C, Messer HH. Micro-tensile bond strengths of bonding agents to pulpal floor dentine. Int Endod J. Oct; 35(10):833-9, 2002.